

# R 语言基础：练习(二)

## Table of Contents

1. 数组向量矩阵下标练习 .....	2
Exercise 1 .....	2
Exercise 2 .....	2
Exercise 3 .....	2
Exercise 4 .....	2
Exercise 5 .....	2
Exercise 6 .....	2
Exercise 7 .....	2
Exercise 8 .....	3
Exercise 9 .....	3
Exercise 10 .....	3
2. 因子练习题 .....	3
Exercise 1 .....	3
Exercise 2 .....	3
Exercise 3 .....	3
Exercise 4 .....	4
Exercise 5 .....	4
Exercise 6 .....	4
Exercise 7 .....	4
Exercise 8 .....	4
Exercise 9 .....	5
Exercise 10 .....	5

## 1.数组向量矩阵下标练习

### Exercise 1

If `x <- c("ww", "ee", "ff", "uu", "kk")`, what will be the output for `x[c(2,3)]`?

- a. "ee", "ff"
- b. "ee"
- c. "ff"

### Exercise 2

If `x <- c("ss", "aa", "ff", "kk", "bb")`, what will be the third value in the index vector operation `x[c(2, 4, 4)]`?

- a. "uu"
- b. NA
- c. "kk"

### Exercise 3

If `x <- c("pp", "aa", "gg", "kk", "bb")`, what will be the fourth value in the index vector operation `x[-2]`?

- a. "aa"
- b. "gg"
- c. "bb"

### Exercise 4

Let `a <- c(2, 4, 6, 8)` and `b <- c(TRUE, FALSE, TRUE, FALSE)`, what will be the output for the R expression `max(a[b])`?

### Exercise 5

Let `a <- c(3, 4, 7, 8)` and `b <- c(TRUE, TRUE, FALSE, FALSE)`, what will be the output for the R expression `sum(a[b])`?

### Exercise 6

Write an R expression that will return the sum value of 10 for the vector `x <- c(2, 1, 4, 2, 1, NA)`

### Exercise 7

If `x <- c(1, 3, 5, 7, NA)` write an r expression that will return the output 1, 3, 5, 7.

## Exercise 8

Consider the data frame `s <- data.frame(first= as.factor(c("x", "y", "a", "b", "x", "z")), second=c(2, 4, 6, 8, 10, 12))`. Write an R statement that will return the output 2, 4, 10, by using the variable `first` as an index vector.

## Exercise 9

What will be the output for the R expression `(c(FALSE, TRUE)) || (c(TRUE, TRUE))`?

## Exercise 10

Write an R expression that will return the positions of 3 and 7 in the vector `x <- c(1, 3, 6, 7, 3, 7, 8, 9, 3, 7, 2)`.

## 2. 因子练习题

### Exercise 1

If `x = c(1, 2, 3, 3, 5, 3, 2, 4, NA)`, what are the levels of `factor(x)`?

- 1, 2, 3, 4, 5
- NA
- 1, 2, 3, 4, 5, NA

### Exercise 2

Let `x <- c(11, 22, 47, 47, 11, 47, 11)`. If an R expression `factor(x, levels=c(11, 22, 47), ordered=TRUE)` is executed, what will be the 4th element in the output?

- 11
- 22
- 47

### Exercise 3

If `z <- c("p", "a", "g", "t", "b")`, then which of the following R expressions will replace the third element in `z` with "b".

- `factor(z[3]) <- "b"`
- `levels(z[3]) <- "b"`
- `z[3] <- "b"`

#### Exercise 4

If `z <- factor(c("p", "q", "p", "r", "q"))` and levels of `z` are "p", "q", "r", write an R expression that will change the level "p" to "w" so that `z` is equal to: "w", "q", "w", "r", "q".

#### Exercise 5

If: `s1 <- factor(sample(letters, size=5, replace=TRUE))` and `s2 <- factor(sample(letters, size=5, replace=TRUE))`, write an R expression that will concatenate `s1` and `s2` in a single factor with 10 elements.

#### Exercise 6

Consider the iris data set in R. Write an R expression that will 'cut' the `Sepal.Length` variable and create the following factor with five levels.

```
(4.3, 5.02] (5.02, 5.74] (5.74, 6.46] (6.46, 7.18] (7.18, 7.9]
32 41 42 24 11
```

#### Exercise 7

Consider again the iris data set. Write an R expression that will generate a two-way frequency table with two rows and three columns. The rows should relate to `Sepal.length` (less than 5: TRUE or FALSE) and columns to `Species`, with the following output:

```
setosa versicolor virginica
FALSE 30 49 49
TRUE 20 1 1
```

#### Exercise 8

Consider the factor `responses <- factor(c("Agree", "Agree", "Strongly Agree", "Disagree", "Agree"))`, with the following output:

```
[1] Agree Agree Strongly Agree Disagree Agree
Levels: Agree Disagree Strongly Agree
```

Later it was found that new a level "Strongly Disagree" exists. Write an R expression that will include "strongly disagree" as new level attribute of the factor and returns the following output:

```
[1] Agree Agree Strongly Agree Disagree Agree
Levels: Strongly Agree Agree Disagree Strongly Disagree
```

### Exercise 9

Let `x <- data.frame(q=c(2, 4, 6), p=c("a", "b", "c"))`. Write an R statement that will replace levels a, b, c with labels "fertiliser1", "fertiliser2", "fertiliser3".

### Exercise 10

If `x <- factor(c("high", "low", "medium", "high", "high", "low", "medium"))`, write an R expression that will provide unique numeric values for various levels of x with the following output:

```
levels value 1 high 1 2 low 2 3 medium 3
```

[返回 课程主页](#)。